

Risk Factors for Death in Homeless Adults in Boston

Stephen W. Hwang, MD, MPH; Joan M. Lebow, MD; Michael F. Bierer, MD, MPH; James J. O'Connell, MD; E. John Orav, PhD; Troyen A. Brennan, MD, JD, MPH

Background: Homeless individuals experience high mortality rates. Males, whites, and substance abusers are more likely to die, but other high-risk characteristics are unknown.

Objective: To identify demographic and clinical factors associated with an increased risk of death in homeless individuals.

Methods: We conducted a case-control study of 558 adults who were seen by a health care program for the homeless in Boston, Mass, and who died in 1988 to 1993. Age-matched paired controls were selected from among individuals seen by the program who were alive at the end of 1993. Predictive data were obtained by blinded review of medical records. Odds ratios (ORs) for death were calculated using logistic regression analysis models.

Results: In a multivariate analysis, the strongest risk fac-

tors for death were acquired immunodeficiency syndrome (OR, 55.8), symptomatic human immunodeficiency virus infection (OR, 17.7), asymptomatic human immunodeficiency virus infection (OR, 4.1), renal disease (OR, 18.4), a history of cold-related injury (OR, 8.0), liver disease (OR, 3.8), and arrhythmia (OR, 3.3). A history of substance abuse involving injection drugs (OR, 1.6) or alcohol (OR, 1.5) also increased the risk of mortality. Nonfluency in English was associated with a decreased risk of death (OR, 0.4).

Conclusions: In a group of adults seen by a health care program for the homeless, specific medical illnesses were associated with the greatest risk of death. Substance abuse alone was less strongly associated with death. Interventions to reduce mortality among the homeless should focus on individuals with high-risk characteristics.

Arch Intern Med. 1998;158:1454-1460

From the Inner City Health Program, St Michael's Hospital, Toronto, Ontario (Dr Hwang); Boston Health Care for the Homeless Program (Drs Lebow, Bierer, and O'Connell); the Division of General Medicine, Clinical Initiatives Development Program, Brigham and Women's Hospital (Drs Orav and Brennan), and the Department of Health Policy and Management, Harvard School of Public Health (Dr Brennan), Boston, Mass.

HOMELESSNESS IS a major social problem that affects at least half a million Americans.¹⁻³ Not surprisingly, homeless individuals are at greatly increased risk of death. Studies in Atlanta, Ga, and San Francisco, Calif, found that homeless people often died in their fifth decade, with accidents or homicides causing about half of all deaths.^{4,5} The mortality rate among homeless men in Sweden was 4 times higher than expected.⁶ In Philadelphia, Pa, the age-adjusted mortality rate in a cohort of homeless adults was 3.5 times that of the general population; mortality was particularly high among white men and substance abusers.⁷ Homeless men in Boston, Mass, aged 18 to 24 years and 25 to 44 years were 5.9 and 3.0 times more likely to die, respectively, than their counterparts in the general population.⁸

The prevalence of conditions such as substance abuse, psychiatric disorders, and major medical illnesses among homeless

individuals is high.^{3,9-19} A systematic effort to identify the comorbid conditions that are associated with an increased risk of mortality in the homeless is important, because this information would assist efforts to prevent untimely deaths in this population. High-risk characteristics could be used as screening criteria by clinicians and social service agencies to focus attention on homeless individuals who are most likely to die in the near future. We conducted a case-control study of deaths in a cohort of homeless adults in Boston. Our main objective was to identify risk factors associated with death in this population.

RESULTS

SUBJECTS

A total of 17 292 patients had contact with BHCHP, of whom 606 died in Massachusetts in 1988 to 1993. No medical records could be obtained for 34 decedents; this

SUBJECTS AND METHODS

STUDY POPULATION AND CASE DEFINITION

The study population consisted of adults seen by the Boston Health Care for the Homeless Program (BHCHP) between July 1, 1988, and December 31, 1993. Patients were seen at more than 40 sites, including homeless shelters, drop-in centers, outpatient clinics, a recuperative residential facility for the homeless, and on the street. Case patients were identified by comparing the patient database with the Massachusetts death registry from 1988 to 1993. These methods have been described in a previous report.⁸

SELECTION OF CONTROLS

Each case patient was paired with a control from the patient database who was born in the same year and who was alive at the end of 1993. Case patients and controls were matched for the month and year of first contact with the program, so that their potential periods of observation were equal. By selecting controls from a population of homeless patients, we sought to determine which factors other than homelessness per se were associated with death. Case patients and controls were not matched for sex because we wished to determine the effect of sex on risk of mortality.

If no control matched for year of birth was found, controls born within ± 2 years of the case patient were accepted; if no such controls were found, the case patient was eliminated from the analysis. If 2 or more potential controls were found, 1 was selected randomly. If no medical records could be obtained for a control patient, a replacement was selected randomly from the remaining candidates. To eliminate individuals who died outside of Massachusetts from the control group, controls were matched against the National Death Index for 1988 to 1993.²⁰ Controls who had died out of state were replaced in the same manner as controls without medical records.

DATA COLLECTION

All patient data were obtained by retrospective review of medical records. Anticipating that homeless individuals often receive health care at multiple sites, medical records were sought at BHCHP and at Boston City Hospital, New England Medical Center, and Massachusetts General Hospital. These institutions were selected because they are located close to the largest homeless shelters in Boston and provide the majority of hospital-based care for patients seen by BHCHP.

Medical records were reviewed by physicians who were blinded to the vital status of all patients and who did not review the chart of any patient they knew. Data were abstracted using standardized forms and explicit definitions for comorbidities. Chart review was restricted to records dated no more than 2 months after first contact to identify baseline conditions rather than conditions that developed during follow-up. The 2-month interval was chosen to allow for the reporting of any diagnostic testing initiated at the time of first contact.

Definitions of medical conditions were taken from the Charlson Comorbidity Index,²¹ with the addition of certain conditions believed to be prevalent among homeless individuals (**Table 1**). Injuries and poisonings were categorized according to the *International Classification of Diseases, Ninth Revision, Clinical Modification*.²⁴ If more than 10 injuries or poisonings had occurred, only the 10 most recent events were recorded. Mental illnesses were grouped in 3 broad categories defined in the *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition*²⁵: schizophrenia or other psychotic disorder, mood disorder, and any other psychiatric disorder. Currently prescribed psychotropic medications and any history of a suicide attempt resulting in hospitalization were noted. Any history of or current substance abuse involving alcohol, opiates, cocaine, sedative-hypnotics, or injection drug use was recorded.

STATISTICAL ANALYSIS

Matched-pairs logistic regression analysis models were used to estimate the odds ratio and 95% confidence interval associated with each potential risk factor. First, simple univariate models were constructed; factors that were associated with mortality at the $P \leq .05$ level were candidates for entry into the multivariate model. The multivariate model was built using a stepwise selection procedure with $P \leq .05$ required for both entry and retention. Demographic factors were added to the model if they were believed to be essential. Confounding factors were identified if their inclusion changed the β coefficient of any factor by more than 10%. All analyses were performed using SAS-PC statistical software (SAS Institute Inc, Cary, NC).

Attributable risk was calculated for each factor using the prevalence of the factor in control subjects to estimate the prevalence of the factor in the homeless population in general. The attributable risk estimates the proportion of all deaths that are attributable to a particular factor, assuming those factors that increase the risk of death are causally related to death.²⁶

group did not differ significantly in age or sex from all other decedents. No suitable age-matched control could be identified for 14 older patients (average age, 69.4 years), all of whom were men. These 48 case patients were excluded from subsequent analyses. In the 558 case patients included in the study, the average age at first contact was 44.8 years (range, 19.4-80.5 years). Death occurred an average of 25.2 months after first contact.

Medical records were obtained for 85% of the 558 initially selected control patients. The remaining 15% were replaced by controls whose medical records were obtainable. Matching of the 558 control patients against the Na-

tional Death Index revealed that 9 had died out of state; these patients were replaced with alternate controls.

The demographic characteristics of case patients and controls are shown in **Table 2**. Significantly fewer deaths occurred among women and individuals of Hispanic race. Nonfluency in English was associated with a more than 3-fold decrease in the risk of death.

MEDICAL CONDITIONS

Certain medical conditions were associated with a greatly increased risk of death in univariate analyses (**Table 3**).

Table 1. Definitions of Conditions

Condition	Definition
Nonfluency in English	Inability to communicate with health care providers in English without the use of an interpreter
Thromboembolic disease	Deep vein thrombosis diagnosed using ultrasonography or venography, or pulmonary embolism diagnosed using ventilation-perfusion scan or pulmonary angiography
Chronic venous insufficiency	Lower extremity edema attributed to venous insufficiency, with characteristic skin changes and/or stasis ulceration
Active tuberculosis	Definitive diagnosis based on positive cultures for <i>Mycobacterium tuberculosis</i> , or presumptive diagnosis followed by treatment with a course of multidrug therapy
Endocarditis	Definitive diagnosis of endocarditis by Duke ²² criteria
Human immunodeficiency virus (HIV) infection, asymptomatic	Clinical category A according to the Centers for Disease Control and Prevention (CDC) ²³ 1993 revised classification system for HIV infection
HIV infection, symptomatic	Clinical category B according to the CDC 1993 revised classification system for HIV infection
HIV infection, acquired immunodeficiency syndrome	Clinical category C according to the CDC 1993 revised classification system for HIV infection
Pancreatitis	Clinical diagnosis of pancreatitis, based on at least 2 of the following: abdominal pain, elevated serum amylase and/or lipase level, radiographic evidence of pancreatitis
Seizures	Any seizure during lifetime
Chronic cognitive impairment	Mental retardation or dementia
Internal injury of the chest, abdomen, or pelvis	Traumatic pneumothorax or hemothorax, or injury to any intrathoracic, intra-abdominal, or pelvic organ
Open wound	Penetrating injuries or traumatic amputations, not involving internal injury as defined above
Minor injury	Nonpenetrating injuries such as superficial lacerations, abrasions, ecchymoses, and contusions
Poisoning or drug overdose	Toxic effects of drugs, medicinal substances, biological substances, or nonmedical substances (excluding acute alcohol intoxication or simple drunkenness)

Human immunodeficiency virus (HIV) infection imparted a degree of risk that varied according to the clinical stage of the disease. A diagnosis of solid tumor with metastases was uniformly associated with death. The risk of mortality was increased more than 6-fold among individuals with thromboembolic disease, liver disease (defined as cirrhosis with or without clinical evidence of portal hypertension), or renal disease (defined as end-stage renal disease or a serum creatinine level of ≥ 177 mmol/L [≥ 2.0 mg/dL]). A high risk of mortality was observed in all individuals with elevated creatinine levels, and not only in those who were dependent on dialysis.

Cardiovascular conditions significantly associated with mortality included congestive heart failure and ar-

Table 2. Demographic Characteristics of 558 Homeless Adults Who Died and Their Matched Controls, and Odds Ratios for Death in Univariate Analyses

Variable	Case Patients, No. (%)	Controls, No. (%)	Odds Ratio (95% CI)*
Sex			
Male†	478 (86)	428 (77)	...
Female	80 (14)	130 (23)	0.6 (0.4-0.8)
Race			
White†	330 (59)	296 (53)	...
Black	169 (30)	177 (32)	0.8 (0.6-1.1)
Hispanic	45 (8)	66 (12)	0.6 (0.4-0.9)
Other	14 (3)	19 (3)	0.7 (0.3-1.4)
Veteran status			
Not a veteran†	479 (86)	505 (91)	...
Veteran	79 (14)	53 (10)	1.7 (1.1-2.6)
English fluency‡			
Fluent†	544 (97)	508 (91)	...
Nonfluent	14 (3)	50 (9)	0.3 (0.2-0.5)

*Odds ratios are calculated as the risk of death among subjects in the demographic category in question, compared with subjects in the reference category. CI indicates confidence interval; ellipses, not applicable.

†Reference category.

‡See Table 1 for definition.

rhythmias. The presence of pulmonary disease (primarily asthma and chronic obstructive pulmonary disease) increased the risk of death to a lesser degree. Other medical conditions that significantly increased the risk of death included a history of seizures, peptic ulcer disease, gastrointestinal bleeding from causes other than peptic ulcer disease, and pancreatitis.

PSYCHIATRIC CONDITIONS AND SUBSTANCE ABUSE

Psychiatric conditions and substance abuse histories are shown in **Table 4**. Among psychiatric conditions, the only significant predictor of death was a history of a suicide attempt. The risk of dying was about twice as high among individuals with a history of abusing alcohol or cocaine. Death was more than 3 times more likely in individuals with a history of opiate abuse or injection drug use.

INJURIES AND POISONINGS

A history of injury or poisoning severe enough to require medical attention was commonly found (**Table 5**). A previous cold-related injury (frostbite, hypothermia, or immersion foot) was associated with an almost 7-fold increase in the risk of death. Death was also more likely to occur in individuals who had previously suffered a fracture or dislocation. A history of poisoning or drug overdose, typically involving opiates or multiple drugs, increased the risk of death almost 3-fold. The external cause of previous injuries was not strongly associated with mortality. The risk of death increased as the number of previous injuries and poisonings increased; death was 1.7 times more likely in individuals who had suffered 4 or more events compared with those who had not.

Table 3. Medical Conditions in Case Patients and Controls, and Odds Ratios for Death in Univariate Analyses*

Variable†	Case Patients, No. (%)	Controls, No. (%)	Odds Ratio (95% CI)
Cardiovascular conditions			
Hypertension	116 (21)	106 (19)	1.1 (0.8-1.5)
Angina	16 (3)	18 (3)	0.9 (0.4-1.8)
Myocardial infarction	23 (4)	16 (3)	1.4 (0.8-2.7)
Congestive heart failure	20 (4)	7 (1)	3.2 (1.3-7.9)
Arrhythmia	26 (5)	6 (1)	5.0 (1.9-13.1)
Valvular heart disease	9 (2)	6 (1)	1.5 (0.5-4.2)
Peripheral vascular disease	11 (2)	5 (1)	2.2 (0.8-6.3)
Thromboembolic disease	20 (4)	3 (1)	6.7 (2.0-22.4)
Chronic venous insufficiency	29 (5)	25 (5)	1.2 (0.7-2.1)
Pulmonary disease	110 (20)	79 (14)	1.5 (1.1-2.1)
Infectious diseases			
Positive tuberculin skin test	94 (17)	90 (16)	1.1 (0.8-1.5)
Active tuberculosis	21 (4)	11 (2)	2.1 (1.0-4.7)
Endocarditis	15 (3)	7 (1)	2.1 (0.9-5.3)
HIV infection, asymptomatic	20 (4)	9 (2)	2.2 (1.0-4.9)
HIV infection, symptomatic	44 (8)	6 (1)	7.3 (3.1-17.2)
AIDS	65 (12)	5 (1)	21.0 (6.6-66.9)
Gastrointestinal conditions			
Peptic ulcer disease	62 (11)	32 (6)	2.2 (1.4-3.4)
Gastrointestinal bleeding	23 (4)	7 (1)	3.3 (1.4-7.7)
Pancreatitis	42 (8)	17 (3)	2.5 (1.4-4.3)
Liver disease	31 (6)	5 (1)	6.2 (2.4-15.9)
Neurologic conditions			
Seizures	133 (24)	71 (13)	2.2 (1.6-3.0)
Chronic cognitive impairment	19 (3)	13 (2)	1.5 (0.7-3.0)
Cerebrovascular disease	10 (2)	10 (2)	1.0 (0.4-2.4)
Cancers			
Solid tumors without metastases	20 (4)	17 (3)	1.2 (0.6-2.5)
Solid tumors with metastases	15 (3)	0 (0)	... ‡
Diabetes mellitus	36 (6)	25 (4)	1.4 (0.9-2.4)
Renal disease	13 (2)	2 (0.4)	6.5 (1.5-28.8)
Rheumatologic disease	5 (1)	2 (0.4)	2.5 (0.5-12.9)

*Odds ratios are calculated as the risk of death among subjects with the condition in question, compared with subjects without it. CI indicates confidence interval; HIV, human immunodeficiency virus; and AIDS, acquired immunodeficiency syndrome.

†See Table 1 for definitions of medical conditions not described in the Charlson Comorbidity Index.

‡Odds ratio is undefined.

RESULTS OF MULTIVARIATE ANALYSES

The results of multivariate analyses are shown in **Table 6**. The strongest predictor of death was the clinical diagnosis of the acquired immunodeficiency syndrome, which was associated with an odds ratio of 55.8 and an attributable risk of 0.35. Symptomatic and asymptomatic HIV infection also significantly increased the risk of mortality. Other medical diagnoses associated with death were renal disease, liver disease, arrhythmia, and seizures. A history of solid tumor with metastases could not be included in the final model, because the odds ratio was undefined. Substance abuse was not as strongly associated

Table 4. Psychiatric History and Substance Abuse in Case Patients and Controls, and Odds Ratios for Death in Univariate Analyses

Variable	Case Patients, No. (%)	Controls, No. (%)	Odds Ratio (95% CI)
Psychiatric history			
Schizophrenia or other psychotic disorder	48 (9)	67 (12)	0.7 (0.5-1.0)
Mood disorder	81 (15)	81 (15)	1.0 (0.7-1.4)
Any other psychiatric diagnosis	55 (10)	52 (9)	1.1 (0.7-1.6)
Previous suicide attempt	47 (8)	22 (4)	2.3 (1.4-4.0)
Psychotropic medications†			
Antipsychotics	27 (5)	42 (8)	0.6 (0.4-1.0)
Antidepressants	17 (3)	13 (2)	1.3 (0.6-2.7)
Antimaniacs	6 (1)	8 (1)	0.8 (0.3-2.2)
Anxiolytics	21 (4)	23 (4)	0.9 (0.5-1.7)
Any psychotropic medication	57 (10)	68 (12)	0.8 (0.6-1.2)
Substance abuse history‡			
Alcohol	426 (89)	339 (79)	2.3 (1.7-3.0)
Opiates	141 (25)	62 (11)	3.1 (2.1-4.4)
Cocaine	139 (25)	89 (16)	1.9 (1.4-2.6)
Sedatives or hypnotics	33 (6)	18 (3)	1.9 (1.1-3.5)
Injection drug use	206 (37)	98 (18)	3.2 (2.3-4.4)

*Odds ratios are calculated as the risk of death among subjects with the condition in question compared with subjects without it. CI indicates confidence interval.

†Refers to psychotropic medications that were currently prescribed for the subject at the time of first contact.

‡Refers to a history of current substance abuse.

with death as were medical diagnoses; the risk of death was increased less than 2-fold in the presence of a history of injection drug use or alcoholism.

A history of either cold-related injury or fracture was a significant risk factor; psychiatric conditions, however, did not meet criteria for inclusion. The trend toward decreased risk of death among women ($P = .08$) and Hispanics ($P = .08$) did not reach statistical significance; sex and race were nonetheless included in the model because of their clinical relevance. Nonfluency in English was associated with a significantly decreased risk of death.

COMMENT

This study elucidates the risk factors for death in a group of homeless adults in Boston. Our results agree with previous findings that male sex, white race, and a history of substance abuse increase the risk of death in homeless individuals.⁷ These criteria, however, are of limited use in identifying a high-risk subgroup because such a large proportion of the homeless population in the United States meets this profile. More than three quarters of the homeless are men, about half are white, and a large number abuse alcohol or cocaine.³ Our study shows that a small subgroup of homeless individuals at extremely high risk of death can be identified by the presence of medical conditions such as HIV infection, renal disease, liver disease, and arrhythmias. These comorbidities are associated with a much greater risk of death than substance abuse alone.

Table 5. Injury and Poisoning Histories in Case Patients and Controls, and Odds Ratios for Death in Univariate Analyses*

Variable	Case Patients, No. (%)	Controls, No. (%)	Odds Ratio (95% CI)
Type of injury			
Fracture	163 (29)	117 (21)	1.6 (1.2-2.2)
Dislocation	11 (2)	2 (0.4)	5.5 (1.2-24.8)
Major head injuries			
Concussion	31 (6)	23 (4)	1.4 (0.8-2.3)
Cerebral contusion	4 (1)	5 (1)	0.8 (0.2-3.0)
Subdural hematoma	11 (2)	6 (1)	1.8 (0.7-5.0)
Intracerebral hemorrhage	4 (1)	3 (1)	1.3 (0.3-6.0)
Any major head injury	47 (4)	34 (3)	1.4 (0.9-2.2)
Internal injury of the chest, abdomen, or pelvis†	30 (5)	26 (5)	1.2 (0.7-2.0)
Open wound†	33 (6)	27 (5)	1.2 (0.7-2.1)
Frostbite, hypothermia, or immersion foot	21 (4)	4 (1)	6.7 (2.0-22.4)
Poisoning or drug overdose†	31 (6)	12 (2)	2.8 (1.4-5.6)
Burns	25 (4)	26 (5)	1.0 (0.6-1.7)
Minor injury†	204 (37)	186 (33)	1.2 (0.9-1.5)
External cause of injury			
Assault			
Without knife or gun	175 (31)	143 (26)	1.3 (1.0-1.7)
With knife	43 (8)	46 (8)	0.9 (0.6-1.4)
With gun	17 (3)	14 (3)	1.2 (0.6-2.5)
Any assault	204 (37)	169 (30)	1.3 (1.0-1.7)
Motor vehicle crash	42 (8)	42 (8)	1.0 (0.6-1.6)
Pedestrian-motor vehicle crash	26 (5)	24 (4)	1.1 (0.6-1.9)
Accident (not involving motor vehicle)	145 (26)	107 (19)	1.5 (1.1-2.0)
No. of injuries and poisonings			
≤3‡	471 (84)	506 (91)	...
≥4	87 (16)	52 (9)	1.7 (1.2-2.5)

*Odds ratios are calculated as the risk of death among subjects with the condition in question, compared with subjects without it. CI indicates confidence interval; ellipses, not applicable.

†See Table 1 for definitions.

‡Reference category.

Despite these findings, substance abuse remains an important risk factor for death for 2 reasons. First, conditions that moderately increase risk of death will be linked to a large number of deaths if the condition is common. Although the odds ratio associated with alcoholism was only 1.5, the high prevalence of alcoholism in the homeless population led to a large attributable risk. By this measure, more than a quarter of deaths in this cohort were attributable to alcohol abuse.

Second, conditions such as liver disease, seizures, cold-related injuries, and fractures may in fact be surrogate markers for severity of substance abuse. For example, a history of cold-related injury greatly increased the risk of death, but only 4 individuals actually died of hypothermia.⁸ A history of frostbite or hypothermia may in fact identify a group of patients with severe substance abuse who demonstrate a high level of self-neglect and risk-taking behavior. These characteristics may predispose individuals to death from a variety of causes. Substance abuse is likely a major contributing factor to deaths because of injuries and drug overdoses; in this cohort,

13% of all deaths were accidental, 4% were caused by homicide, and 2% were caused by suicide.⁸

Clinical interventions to prevent deaths among the homeless may be most successful when focused on high-risk subgroups such as individuals with HIV infection. Acquired immunodeficiency syndrome was the leading cause of death in our cohort, accounting for 18% of all deaths.⁸ Since newer treatments for HIV infection can reduce morbidity and prolong life,²⁷ ensuring that HIV-infected homeless individuals have access to appropriate medical care is of greater importance than ever before.²⁸ At a social policy level, efforts should be redoubled to prevent HIV infection among homeless individuals who engage in high-risk behaviors. In our study, 75% of those who died of acquired immunodeficiency syndrome had a history of injection drug use. Methadone maintenance treatment²⁹⁻³¹ and needle-exchange programs^{32,33} are effective methods of reducing HIV infection in injection drug users and should be expanded.

Homeless individuals with conditions such as renal disease, liver disease, arrhythmias, and seizures have a markedly increased risk of death. Primary care providers who are experienced in working with homeless patients should orchestrate clinical care for these individuals. Intensive efforts to engage patients, encourage adherence to therapy, and trace patients who are lost to follow-up may be most effectively coordinated through multidisciplinary teams of physicians, nurses, and social workers.³⁴ Treatment of coexisting substance abuse should also be emphasized.

The 3-fold decrease in risk of death associated with nonfluency in English is an interesting and unexpected finding. This correlation cannot be explained completely on the basis of confounding by Hispanic race, since it remains significant even in a multivariate model that includes race as a predictive factor. Nonfluency in English may identify a subgroup of individuals who have become homeless because of purely economic reasons and who therefore suffer from a lower burden of disease than do other homeless people. In other words, individuals who are fluent in English may be less likely to become homeless unless, in addition to being poor, they have comorbid conditions that predispose them to both homelessness and death.

We found a trend toward a decreased risk of death among individuals diagnosed as having schizophrenia or other psychotic disorders. In Massachusetts, individuals with these conditions are eligible for long-term psychiatric case-management services that are coordinated with general medical care. We hypothesize that the paradoxically beneficial effect of schizophrenia may be the result of the structured medical and social services that are provided to these chronically disabled patients. Therefore, at a health policy level, expanding the availability of these types of supportive services might reduce deaths among homeless individuals.

Certain limitations of this study should be noted. Only deaths that occurred in Massachusetts were identified; risk factors in individuals who migrated out of state and then died were not examined. We studied individuals who had contact with a health care program for the homeless, and thus our findings may not be generaliz-

Table 6. Risk Factors Associated With Death in Homeless Adults in Multivariate Analyses*

Variable†	Case Patients, No. (%)	Controls, No. (%)	Odds Ratio (95% CI)	Attributable Risk‡
HIV infection, AIDS	65 (12)	5 (1)	55.8 (14.4-216.9)	0.35
HIV infection, symptomatic	44 (8)	6 (1)	17.7 (5.3-58.8)	0.14
HIV infection, asymptomatic	20 (4)	9 (2)	4.1 (1.5-11.6)	0.06
Renal disease	13 (2)	2 (0.4)	18.4 (2.1-161.5)	0.07
Frostbite, hypothermia, or immersion foot	21 (4)	4 (1)	8.0 (1.7-38.6)	0.07
Liver disease§	31 (6)	5 (1)	3.8 (1.2-11.5)	0.03
Arrhythmia	26 (5)	6 (1)	3.3 (1.1-9.4)	0.02
Seizures§	133 (24)	71 (13)	1.7 (1.1-2.6)	0.08
Fracture	163 (29)	117 (21)	1.6 (1.1-2.3)	0.11
Injection drug use§	206 (37)	98 (18)	1.6 (1.1-2.3)	0.10
Alcohol dependence or abuse	426 (89)	339 (79)	1.5 (1.1-2.3)	0.28
Nonfluency in English	14 (3)	50 (9)	0.4 (0.2-0.8)	...
Female	80 (14)	130 (23)	0.7 (0.5-1.1)	...
Black¶	169 (30)	177 (32)	0.9 (0.6-1.2)	...
Hispanic¶	45 (8)	66 (12)	0.6 (0.3-1.1)	...
Other race¶	14 (3)	19 (3)	0.7 (0.3-1.5)	...

*Odds ratios are calculated as the risk of death among subjects with the characteristic in question, compared with subjects without it, unless otherwise specified. CI indicates confidence interval; HIV, human immunodeficiency virus; AIDS, acquired immunodeficiency syndrome; and ellipses, not applicable.

†See Table 1 for definitions.

‡Attributable risk is an estimate of the proportion of all deaths that are attributable to a particular risk factor, assuming that factors that increase risk are causally related to death.

§Positive confounding existed between the following pairs of factors: liver disease and gastrointestinal bleeding, liver disease and pancreatitis, seizures and pancreatitis, and injection drug use and previous suicide attempt. If these confounding factors were added to the multivariate model, the odds ratio (95% CI) for liver disease became 2.9 (1.0-9.0), for seizures 1.5 (1.0-2.4), and for injection drug use 1.5 (1.0-2.3).

||Reference category is male.

¶Reference category is white.

able to homeless people who do not seek health care at all. Previous investigators³⁵ have found that homeless people who seek health care tend to have higher levels of morbidity than non-care seekers. While all the individuals in our study were homeless at the time of first contact, data are not available on which individuals subsequently obtained temporary or permanent housing. Thus, our analysis did not take the subsequent duration of homelessness into account.

Characteristics of case patients and controls were obtained by retrospective chart review. This method may underestimate the prevalence of comorbidities in homeless individuals who have not undergone a comprehensive health evaluation.³⁶ Thus, the association between certain risk factors and death may have been overestimated if case patients tended to receive more medical scrutiny than controls prior to entry into the study. On the other hand, 15% of initial controls were replaced because they had no medical records. This process preferentially selected controls with more contact with the health care system and may have weakened true associations between risk factors and death.

In conclusion, our study identifies the characteristics of homeless individuals at extremely high risk of death. Clinicians who see homeless individuals should be alert to the need for aggressive intervention and close medical follow-up in patients with these risk factors. To prevent many of these deaths, however, changes must occur at the level of social policy rather than at the level of individual clinical intervention. These changes include wider availability of supportive services and stable low-income housing, improved access to health care, and the expansion of methadone maintenance treatment and

needle-exchange programs. Efforts to prevent deaths among the homeless must address the complex medical, psychiatric, substance abuse, and socioeconomic problems of this vulnerable population. Studies are needed to rigorously evaluate the effectiveness of these efforts.

Accepted for publication November 13, 1997.

This study was supported in part by a grant from the Department of Medicine, Boston Medical Center, Boston, Mass. Dr Hwang is a recipient of a Health Services Research Fellowship from the Agency for Health Care Policy and Research, Rockville, Md.

We thank Alice Knowles of JSI Incorporated for invaluable data management services and Charlene Zion of the Massachusetts Registry of Vital Records and Statistics for providing death registry data (Boston).

Reprints: Stephen W. Hwang, MD, MPH, Inner City Health Program, St Michael's Hospital, Victoria Wing 4-154, 30 Bond St, Toronto, Ontario, Canada M5B 1W8 (e-mail: hwangs@smh.toronto.on.ca).

REFERENCES

- Hombs ME, Snyder M. *Homelessness in America: A Forced March to Nowhere*. Washington, DC: Community for Creative Non-Violence; 1982.
- Burt M, Cohen B. *America's Homeless: Numbers, Characteristics, and Programs That Serve Them*. Washington, DC: The Urban Institute Press; 1989.
- Jencks C. *The Homeless*. Cambridge, Mass: Harvard University Press; 1994.
- Deaths among the homeless: Atlanta, Georgia. *MMWR Morb Mortal Wkly Rep*. 1987;36:297-299.
- Deaths among homeless persons: San Francisco, 1985-1990. *MMWR Morb Mortal Wkly Rep*. 1991;40:877-880.
- Alstrom CH, Lindelius R, Salum I. Mortality among homeless men. *Br J Addict*. 1975;70:245-252.

7. Hibbs JR, Benner L, Klugman L, et al. Mortality in a cohort of homeless adults in Philadelphia. *N Engl J Med*. 1994;331:304-309.
8. Hwang SW, Orav EJ, O'Connell JJ, Lebow JM, Brennan TA. Causes of death in homeless adults in Boston. *Ann Intern Med*. 1997;126:625-628.
9. Bassuk EL, Rubin L, Lauriat A. Is homelessness a mental health problem? *Am J Psychiatry*. 1984;141:1546-1550.
10. Koegel P, Burnam A, Farr RK. The prevalence of specific psychiatric disorders among homeless individuals in the inner city of Los Angeles. *Arch Gen Psychiatry*. 1988;45:1085-1092.
11. Susser E, Struening EL, Conomver S. Psychiatric problems in homeless men: lifetime psychosis, substance use, and current distress in new arrivals at New York City shelters. *Arch Gen Psychiatry*. 1989;46:845-850.
12. Breakey WR, Rischer PJ, Kramer M, et al. Health and mental health problems of homeless men and women in Baltimore. *JAMA*. 1989;262:1352-1357.
13. Ferenchick GS. Medical problems of homeless and nonhomeless persons attending an inner-city health clinic: a comparative study. *Am J Med Sci*. 1991;301:379-382.
14. Characteristics and risk behaviors of homeless black men seeking services from the Community Homeless Assistance Plan: Dade County, Florida, August 1991. *MMWR Morb Mortal Wkly Rep*. 1991;40:865-868.
15. Torres RA, Mani S, Altholz J, Brickner PW. Human immunodeficiency virus infection among homeless men in a New York City shelter. *Arch Intern Med*. 1990;150:2030-2035.
16. Brudney K, Dobkin J. Resurgent tuberculosis in New York City: human immunodeficiency virus, homelessness, and the decline of tuberculosis control programs. *Am Rev Respir Dis*. 1991;144:745-749.
17. Allen DM, Lehman JS, Green TA, et al. HIV infection among homeless adults and runaway youth, United States, 1989-1992. *AIDS*. 1994;8:1593-1598.
18. Zolopa AR, Hahn JA, Gorter R, et al. HIV and tuberculosis infection in San Francisco's homeless adults: prevalence and risk factors in a representative sample. *JAMA*. 1994;272:455-461.
19. Concato J, Rom WN. Endemic tuberculosis among homeless men in New York City. *Arch Intern Med*. 1994;154:2069-2073.
20. *National Death Index User's Manual*. Hyattsville, Md: National Center for Health Statistics; 1990. US Dept of Health and Human Services publication PHS 90-1148.
21. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40:373-383.
22. *International Classification of Diseases, Ninth Revision, Clinical Modification*. 5th ed. Salt Lake City, Utah: Medicode; 1996.
23. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition*. Washington, DC: American Psychiatric Association; 1994.
24. Kleinbaum DG, Kupper LL, Morgenstern H. *Epidemiologic Research: Principles and Quantitative Methods*. Belmont, Calif: Lifetime Learning; 1982.
25. Carpenter CCJ, Fischl MA, Hammer SM, et al. Antiretroviral therapy for HIV infection in 1997. *JAMA*. 1997;277:1962-1969.
26. Bangsberg D, Tulsy FP, Hecht FM, Moss AR. Protease inhibitors in the homeless. *JAMA*. 1997;278:63-65.
27. Serpelloni G, Carrieri MP, Rezza G, Morganti S, Gomma M, Binkin N. Methadone treatment as a determinant of HIV risk reduction among injecting drug users: a nested case-control study. *AIDS Care*. 1994;6:215-220.
28. Metzger DS, Woody GE, McLellan AT, O'Brien CP, Druley P, Navaline H. Human immunodeficiency virus seroconversion among intravenous drug users in- and out-of-treatment: an 18-month prospective follow-up. *J Acquir Immune Defic Syndr Hum Retrovirol*. 1993;6:1049-1056.
29. Williams AB, McNella EA, Williams AE, D'Aquila RT. Methadone maintenance treatment and HIV type 1 seroconversion among injecting drug users. *AIDS Care*. 1992;4:35-41.
30. Kaplan EH. A method for evaluating needle exchange programmes. *Stat Med*. 1994;13:2179-2187.
31. Kaplan EH, Heimer R. HIV incidence among New Haven needle exchange participants: updated estimates from syringe tracking and testing data. *J Acquir Immune Defic Syndr Hum Retrovirol*. 1995;10:175-176.
32. Brickner PW, Scanlan BC, Conanan B, et al. Homeless persons and health care. *Ann Intern Med*. 1986;104:405-409.
33. Padgett D, Struening EL, Andrews H. Factors affecting the use of medical, mental health, alcohol, and drug treatment services by homeless adults. *Med Care*. 1990;28:805-821.
34. Gelberg L, Linn LS. Assessing the physical health of homeless adults. *JAMA*. 1989;262:1973-1979.
35. Durack DT, Lukes AS, Bright DK. New criteria for diagnosis of infective endocarditis: utilization of specific echocardiographic findings: Duke Endocarditis Service. *Am J Med*. 1994;96:200-209.
36. 1993 revised classification system for HIV infection and expanded surveillance case definition for AIDS among adolescents and adults. *MMWR Morb Mortal Wkly Rep*. 1992;41(RR-17):1-19.